

# TETRACOMPOSITION

Liivi Kluge and Eugen Paal

**Abstract.** We consider basic algebraic constructions associated with an *abstract* pre-operad, such as a  $\smile$ -algebra, total composition  $\bullet$ , *pre-coboundary* operator  $\delta$ , *tribraces*  $\{\cdot, \cdot, \cdot\}$  and *tetrabraces*  $\{\cdot, \cdot, \cdot, \cdot\}$ . A derivation deviation of the pre-coboundary operator over the tetrabraces is calculated in terms of the  $\smile$ -multiplication and tribraces.

**Classification (MSC2000).** 18D50.

**Key words.** Comp(osition), (pre-)operad, cup, pre-coboundary, (tri and tetra)braces, derivation deviation.

## 1. INTRODUCTION AND OUTLINE OF THE PAPER

We consider basic algebraic constructions associated with an *abstract* pre-operad  $C$  (Sec. 2), such as a  $\smile$ -algebra (Sec. 3), total composition  $\bullet$  (Sec. 3), *pre-coboundary* operator  $\delta$  (Sec. 4), *tribraces*  $\{\cdot, \cdot, \cdot\}$  (Sec. 5) and *tetrabraces*  $\{\cdot, \cdot, \cdot, \cdot\}$  (Sec. 6). Main result of the present paper is the Main Theorem of Sec. 6. By defining (see degree notations in Sec. 2) a *derivation deviation* (Sec. 6) of the pre-coboundary operator  $\delta$  over the tetrabraces,

$$\begin{aligned} \text{dev}_{\{\cdot, \cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g \otimes b) &:= \delta\{h, f, g, b\} - \{h, f, g, \delta b\} - (-1)^{|b|} \{h, f, \delta g, b\} \\ &\quad - (-1)^{|b|+|g|} \{h, \delta f, g, b\} - (-1)^{|b|+|g|+|f|} \{\delta h, f, g, b\}, \end{aligned}$$

the Main Theorem states that in a the pre-operad  $C$  one has

$$\begin{aligned} (-1)^{|b|} \text{dev}_{\{\cdot, \cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g \otimes b) &= \{h, f, g\} \smile b - \{h, f, g \smile b\} \\ &\quad - (-1)^{|g|} \{h, f \smile g, b\} + (-1)^{|h|f+|g|} f \smile \{h, g, b\}. \end{aligned}$$

Our result is similar to that of [5, 14], where it was announced that for the *Hochschild cochains* the extra terms with  $\delta h$ ,  $\delta f$ ,  $\delta g$  and  $\delta b$  also appear. We work out these extra terms for an *abstract* pre-operad and give their interpretation via a *derivation deviation* of  $\delta$  over the tetrabraces.

The paper can be seen as an an extension of [6] and [7].

## 2. PRE-OPERAD (COMPOSITION SYSTEM)

Let  $K$  be a unital commutative associative ring, and let  $C^n$  ( $n \in \mathbb{N}$ ) be unital  $K$ -modules. For *homogeneous*  $f \in C^n$ , we refer to  $n$  as the *degree* of  $f$  and often write (when it does not cause confusion)  $f$  instead of  $\deg f$ ; for example  $(-1)^f := (-1)^n$ ,  $C^f := C^n$  and  $\circ_f := \circ_n$ . Also, it is convenient to use the shifted (*desuspended*) degree  $|f| := n - 1$ . Throughout this paper, we assume that  $\otimes := \otimes_K$  and work with *homogeneous* elements of  $C$ .

**Definition 2.1.** A linear (right) *pre-operad* (*composition system*) with coefficients in  $K$  is a sequence  $C := \{C^n\}_{n \in \mathbb{N}}$  of unital  $K$ -modules (an  $\mathbb{N}$ -graded  $K$ -module), such that the following conditions hold:

1. For  $0 \leq i \leq m - 1$  there exist *partial compositions*

$$\circ_i \in \text{Hom}(C^m \otimes C^n, C^{m+n-1}), \quad |\circ_i| = 0.$$

2. For all  $h \otimes f \otimes g \in C^h \otimes C^f \otimes C^g$ , the *composition relations* hold,

$$(h \circ_i f) \circ_j g = \begin{cases} (-1)^{|f||g|} (h \circ_j g) \circ_{i+|g|} f & \text{if } 0 \leq j \leq i - 1, \\ h \circ_i (f \circ_{j-i} g) & \text{if } i \leq j \leq i + |f|, \\ (-1)^{|f||g|} (h \circ_{j-|f|} g) \circ_i f & \text{if } i + f \leq j \leq |h| + |f|. \end{cases}$$

3. There exists a unit  $I \in C^1$  such that

$$I \circ_0 f = f = f \circ_i I, \quad 0 \leq i \leq |f|.$$

In the 2nd item, the *first* and *third* parts of the defining relations turn out to be equivalent.

*Remarks 2.2.* A pre-operad is also called a *comp(osition) algebra* or *asymmetric operad* or *non-symmetric operad* or *non- $\Sigma$  operad*. The concept of (*symmetric*) *operad* was formalized by P. May [10] as a tool for the theory of iterated loop spaces. Recent studies and applications can be found in [8].

Above we modified the Gerstenhaber *comp algebra* defining relations [1, 4] by introducing the sign  $(-1)^{|f||g|}$  in the defining relations of the pre-operad. The modification enables us to keep track of (control) sign changes more effectively. One should also note that (up to sign) our  $\circ_i$  is Gerstenhaber's  $\circ_{i+1}$  from [4, 1]; we use the original (non-shifted) convention from [2, 3].

**Example 2.3** (endomorphism pre-operad). Let  $A$  be a unital  $K$ -module and  $\mathcal{E}_A^n := \text{End}_A^n := \text{Hom}(A^{\otimes n}, A)$ . Define (cf. [2, 3, 4]) the partial compositions for  $f \otimes g \in \mathcal{E}_A^f \otimes \mathcal{E}_A^g$  as

$$f \circ_i g := (-1)^{i|g|} f \circ (\text{id}_A^{\otimes i} \otimes g \otimes \text{id}_A^{\otimes (|f|-i)}), \quad 0 \leq i \leq |f|.$$

Then  $\mathcal{E}_A := \{\mathcal{E}_A^n\}_{n \in \mathbb{N}}$  is a pre-operad (with the unit  $\text{id}_A \in \mathcal{E}_A^1$ ) called the *endomorphism pre-operad* of  $A$ . A few examples (without the sign factor) can be found in [3, 4] as well. We use the original indexing of [2, 3] for the defining formulae.

**Example 2.4** (associahedra). A geometrical example of a pre-operad is provided by the Stasheff *associahedra*, which was first constructed in [12]. Quite a surprising realization of the associahedra as *truncated simplices* was discovered and studied in [11, 13, 9].

**Notations 2.5** (scope of a pre-operad). The *scope* of  $(h \circ_i f) \circ_j g$  is given by

$$0 \leq i \leq |h|, \quad 0 \leq j \leq |f| + |h|.$$

It follows from the defining relations of a pre-operad that the scope is a disjoint union of

$$\begin{aligned} B &:= \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid 1 \leq i \leq |h|; 0 \leq j \leq i - 1\}, \\ A &:= \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid 0 \leq i \leq |h|; i \leq j \leq i + |f|\}, \\ G &:= \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid 0 \leq i \leq |h| - 1; i + f \leq j \leq |f| + |h|\}. \end{aligned}$$

Note that the triangles  $B$  and  $G$  are symmetrically situated with respect to the parallelogram  $A$  in the scope  $B \sqcup A \sqcup G$ . The (recommended and impressive) picture is left for a reader as an exercise.

**Recapitulation 2.6.** The defining relations of a pre-operad read

$$(h \circ_i f) \circ_j g = \begin{cases} (-1)^{|f||g|} (h \circ_j g) \circ_{i+|g|} f & \text{if } (i, j) \in B, \\ h \circ_i (f \circ_{j-i} g) & \text{if } (i, j) \in A, \\ (-1)^{|f||g|} (h \circ_{j-|f|} g) \circ_i f & \text{if } (i, j) \in G. \end{cases}$$

The *first* ( $B$ ) and *third* ( $G$ ) parts of the relations are equivalent.

### 3. CUP AND TOTAL COMPOSITION

In Sections 3-5, we recall from [6, 7] main facts about cup-multiplication, total composition, pre-coboundary operator and tribraces in an abstract pre-operad. Fix  $\mu \in C^2$ .

**Definition 3.1.** In a pre-operad  $C$ , the *cup-multiplication*  $\smile := \smile_\mu: C^f \otimes C^g \rightarrow C^{f+g}$  is defined (cf. [2, 3, 4]) by

$$f \smile g := (-1)^f (\mu \circ_0 f) \circ_f g \in C^{f+g}, \quad |\smile| = 1, \quad f \otimes g \in C^f \otimes C^g.$$

Note that  $\circ_f := \circ_{\deg f}$ , and  $|\smile| = 1$  means that  $\deg \smile = 0$ . The pair  $\text{Cup } C := \{C, \smile\}$  is called a  $\smile$ -algebra of  $C$ .

**Example 3.2.** For the endomorphism pre-operad (Example 2.3)  $\mathcal{E}_A$ , one has

$$f \smile g = (-1)^{fg} \mu \circ (f \otimes g) \in \mathcal{E}_A^{f+g}, \quad \mu \otimes f \otimes g \in \mathcal{E}_A^2 \otimes \mathcal{E}_A^f \otimes \mathcal{E}_A^g.$$

**Proposition 3.3.** In a pre-operad  $C$ , one has

$$\mu \circ_0 f = (-1)^f f \smile \text{I}, \quad \mu \circ_1 f = -\text{I} \smile f, \quad f \smile g = -(-1)^{|f|g} (\mu \circ_1 g) \circ_0 f.$$

**Lemma 3.4** ([6, 7]). In a pre-operad  $C$ , the following composition relations hold:

$$(f \smile g) \circ_j h = \begin{cases} (-1)^{g|h|} (f \circ_j h) \smile g & \text{if } 0 \leq j \leq |f|, \\ f \smile (g \circ_{j-f} h) & \text{if } f \leq j \leq |g| + f. \end{cases}$$

**Definition 3.5.** In a pre-operad  $C$ , the *total composition*  $\bullet: C^f \otimes C^g \rightarrow C^{f+g-1}$  is defined (cf. [4, 1]) by

$$f \bullet g := \sum_{i=0}^{|f|} f \circ_i g \in C^{f+g-1}, \quad |\bullet| = 0, \quad f \otimes g \in C^f \otimes C^g.$$

The pair  $\text{Com } C := \{C, \bullet\}$  is called the *composition algebra* of  $C$ .

**Theorem 3.6** ([6, 7]). *In a pre-operad  $C$ , one has*

$$(f \smile g) \bullet h = f \smile (g \bullet h) + (-1)^{|h|g}(f \bullet h) \smile g.$$

*Remark 3.7.* This theorem tells us that *right* translations in  $\text{Com } C$  are (right) derivations of the  $\smile$ -algebra. It turns out that the *left* translations in  $\text{Com } C$  are not derivations of the  $\smile$ -algebra (see Theorem 5.5).

#### 4. PRE-COBOUNDARY OPERATOR

**Definition 4.1.** In a pre-operad  $C$ , define a *pre-coboundary* operator  $\delta := \delta_\mu$  by

$$-\delta f := [f, \mu] := f \bullet \mu - (-1)^{|f|} \mu \bullet f, \quad \mu \otimes f \in C^2 \otimes C^f.$$

**Example 4.2.** In the Gerstenhaber theory [2],  $\delta_\mu$  is the Hochschild *coboundary operator* with the property  $\delta_\mu^2 = 0$ , the latter is due to the *associativity*  $\mu \bullet \mu = 0$ .

In this paper, we do not assume the associativity constraint for  $\mu$ , thus in general  $\delta_\mu^2 \neq 0$ .

**Proposition 4.3.** *In a pre-operad  $C$ , one has*

$$-\delta f = f \smile \text{I} + f \bullet \mu + (-1)^{|f|} \text{I} \smile f.$$

**Definition 4.4.** The *derivation deviation* of  $\delta$  over  $\bullet$  is defined by

$$\text{dev}_\bullet \delta (f \otimes g) := \delta(f \bullet g) - f \bullet \delta g - (-1)^{|g|} \delta f \bullet g.$$

**Theorem 4.5** ([6]). *In a pre-operad  $C$ , one has*

$$(-1)^{|g|} \text{dev}_\bullet \delta (f \otimes g) = f \smile g - (-1)^{fg} g \smile f.$$

#### 5. TRIBRACES

**Definition 5.1** (Gerstenhaber tribraces (cf. [2, 1])). Let  $h \otimes f \otimes g \in C^h \otimes C^f \otimes C^g$ . The Gerstenhaber *tribraces*  $\{\cdot, \cdot, \cdot\}$  are defined as a double sum over the triangle  $G$  by

$$\{h, f, g\} := \sum_{(i,j) \in G} (h \circ_i f) \circ_j g \in C^{h+f+g-2}, \quad |\{\cdot, \cdot, \cdot\}| = 0.$$

**Theorem 5.2** (Getzler identity [6]). *In a pre-operad  $C$ , one has*

$$\begin{aligned} (h, f, g) &:= (h \bullet f) \bullet g - h \bullet (f \bullet g) \\ &= \{h, f, g\} + (-1)^{|f||g|} \{h, g, f\}. \end{aligned}$$

**Theorem 5.3** (Gerstenhaber identity). *In a pre-operad  $C$ , one has*

$$(h, f, g) = (-1)^{|f||g|}(h, g, f).$$

*Proof.* Use the Getzler identity.  $\square$

**Definition 5.4.** The *derivation deviation* of  $\delta$  over the tribraces  $\{\cdot, \cdot, \cdot\}$  is defined by

$$\text{dev}_{\{\cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g) := \delta\{h, f, g\} - \{h, f, \delta g\} - (-1)^{|g|}\{h, \delta f, g\} - (-1)^{|g|+|f|}\{\delta h, f, g\}.$$

**Theorem 5.5** ([7]). *In a pre-operad  $C$ , one has*

$$(-1)^{|g|} \text{dev}_{\{\cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g) = (h \bullet f) \smile g + (-1)^{|h||f|} f \smile (h \bullet g) - h \bullet (f \smile g).$$

**Theorem 5.6.** *In a pre-operad  $C$ , one has*

$$(-1)^{|g|} \text{dev}_{\{\cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g) = [h, f] \smile g + (-1)^{|h||f|} f \smile [h, g] - [h, f \smile g].$$

*Proof.* Combine Theorem 5.5 with Theorem 3.6.  $\square$

## 6. MAIN THEOREM AND GERSTENHABER'S METHOD

In this section, we calculate the derivation deviation of  $\delta$  over tetrabraces for an abstract pre-operad. For this we use the Gerstenhaber (auxiliary variables) method [2, 6].

**Definition 6.1** (ground tetrahedron). The *ground tetrahedron*  $T$  (associated with  $h \otimes f \otimes g \in C^h \otimes C^f \otimes C^g$ ) is defined as a set of points in  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$  with  $\mathbb{N}$ -valued coordinates  $i, j, k$ , which satisfy inequalities

$$\begin{cases} 0 \leq i \leq h-3, \\ i+f \leq j \leq h+f-3, \\ j+g \leq k \leq h+f+g-3. \end{cases}$$

In other words,

$$T := \{(i, j, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} \mid 0 \leq i \leq j-f \leq k-f-g \leq h-3\}.$$

**Definition 6.2** (tetrabraces). Let  $h \otimes f \otimes g \otimes b \in C^h \otimes C^f \otimes C^g \otimes C^b$ . Define the tetrabraces  $\{\cdot, \cdot, \cdot, \cdot\}$  by

$$\{h, f, g, b\} = \sum_{(i,j,k) \in T} ((h \circ_i f) \circ_j g) \circ_k b \in C^{h+f+g+b-3}, \quad |\{\cdot, \cdot, \cdot, \cdot\}| = 0.$$

**Definition 6.3.** The *derivation deviation* of  $\delta$  over the tetrabraces  $\{\cdot, \cdot, \cdot, \cdot\}$  is defined by

$$\begin{aligned} \text{dev}_{\{\cdot, \cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g \otimes b) &:= \delta\{h, f, g, b\} - \{h, f, g, \delta b\} - (-1)^{|b|}\{h, f, \delta g, b\} \\ &\quad - (-1)^{|b|+|g|}\{h, \delta f, g, b\} - (-1)^{|b|+|g|+|f|}\{\delta h, f, g, b\}. \end{aligned}$$

**Main Theorem.** *In a pre-operad  $C$ , one has*

$$\begin{aligned} (-1)^{|b|} \text{dev}_{\{\cdot, \cdot, \cdot, \cdot\}} \delta(h \otimes f \otimes g \otimes b) &= \{h, f, g\} \smile b - \{h, f, g \smile b\} \\ &\quad - (-1)^{|g|}\{h, f \smile g, b\} + (-1)^{|h||f|+|g|} f \smile \{h, g, b\}. \end{aligned}$$

To prove this theorem, we use auxiliary variables.

**Notations 6.4** (auxiliary variables). In a pre-operad  $C$ , for  $(i, j, k) \in T$  define the *auxiliary variables*

$$\begin{aligned}
\Gamma_{i+1,j+1,k+1} &:= -(-1)^{|h|+|f|+|g|+|b|} \mathbf{I} \smile (((h \circ_i f) \circ_j g) \circ_k b) \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{i-1} (((h \circ_s \mu) \circ_{i+1} f) \circ_{j+1} g) \circ_{k+1} b \\
&\quad + (-1)^{|f|+|g|+|b|} ((h \circ_i (\mathbf{I} \smile f)) \circ_{j+1} g) \circ_{k+1} b, \\
\Gamma'_{i+1,j+1,k+1} &:= (-1)^{|g|+|b|} ((h \circ_i (f \smile \mathbf{I})) \circ_{j+1} g) \circ_{k+1} b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=i+1}^{j-f} (((h \circ_s \mu) \circ_i f) \circ_{j+1} g) \circ_{k+1} b \\
&\quad + (-1)^{|g|+|b|} ((h \circ_i f) \circ_j (\mathbf{I} \smile g)) \circ_{k+1} b, \\
\Gamma''_{i+1,j+1,k+1} &:= (-1)^{|b|} ((h \circ_i f) \circ_j (g \smile \mathbf{I})) \circ_{k+1} b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=j-|f|+1}^{k-|f|-g} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_{k+1} b \\
&\quad + (-1)^{|b|} ((h \circ_i f) \circ_j g) \circ_k (\mathbf{I} \smile b), \\
\Gamma'''_{i+1,j+1,k+1} &:= ((h \circ_i f) \circ_j g) \circ_k (b \smile \mathbf{I}) \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=k-|f|-|g|+1}^{|h|} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\
&\quad - (((h \circ_i f) \circ_j g) \circ_k b) \smile \mathbf{I}.
\end{aligned}$$

**Lemma 6.5.** In a pre-operad  $C$ , for  $(i, j, k) \in T$  one has

$$\begin{aligned}
&\delta(((h \circ_i f) \circ_j g) \circ_k b) - ((h \circ_i f) \circ_j g) \circ_k \delta b - (-1)^{|b|} ((h \circ_i f) \circ_j \delta g) \circ_{k+1} b \\
&\quad - (-1)^{|b|+|g|} ((h \circ_i \delta f) \circ_{j+1} g) \circ_{k+1} b \\
&= \Gamma_{i+1,j+1,k+1} + \Gamma'_{i+1,j+1,k+1} + \Gamma''_{i+1,j+1,k+1} + \Gamma'''_{i+1,j+1,k+1}.
\end{aligned}$$

*Proof.* See Appendix A. □

**Notations 6.6** (truncated envelope of  $T'$ ). Now define a *shifted* ground tetrahedron

$$T' := \{(i, j, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} \mid 1 \leq i \leq j - f \leq k - f - g \leq h - 2\}.$$

Its *envelope* is the tetrahedron

$$T'_{env} := \{(i, j, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} \mid 0 \leq i \leq j - |f| \leq k - |f| - |g| \leq h + 1\}.$$

The *boundary* of the envelope  $T'_{env}$  is evidently  $\partial T'_{env} = T'_{env} \setminus T'$ . The *truncated envelope*  $T'_{\widehat{env}}$  of  $T'$  is defined by *removing* (six) edges of  $T'_{env}$ ,

$$\begin{aligned} T'_{\widehat{env}} &:= T'_{env} \setminus \{(i, j, k) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} \mid (0 \leq i \leq h+1, i+|f|, h+f+|g|); \\ &\quad (0, |f|, |f|+|g| \leq k \leq h+f+|g|); \\ &\quad (0 \leq i \leq h+1, i+|f|, i+|f|+|g|); \\ &\quad (0, |f| \leq j \leq h+f, h+f+|g|); \\ &\quad (0 \leq i \leq h+1, h+f, h+f+|g|); \\ &\quad (0, |f| \leq j \leq h+f, j+|g|)\} \\ &= T' \sqcup \partial T'_{\widehat{env}}. \end{aligned}$$

**Lemma 6.7.** *In a pre-operad  $C$ , for  $0 \leq i \leq j-f \leq k-f-g \leq h-2$  one has*

$$(-1)^{|f|+|g|+|b|}((\delta h \circ_i f) \circ_j g) \circ_k b = \Gamma_{ijk} + \Gamma'_{i+1,jk} + \Gamma''_{i+1,j+1,k} + \Gamma'''_{i+1,j+1,k+1},$$

*by definition for  $\Gamma_{0jk}$ ,  $\Gamma'_{i,i+|f|,k}$ ,  $\Gamma''_{ij,j+|g|}$ , and  $\Gamma'''_{ij,|h|+f+g}$  (boundary values on  $\partial T'_{\widehat{env}}$ ).*

*Proof.* See Appendix B.  $\square$

**Boundary Lemma.** *In a pre-operad  $C$ , for  $(i, j, k) \in \partial T'_{\widehat{env}}$  one has*

$$\begin{aligned} \Gamma_{0jk} &= (-1)^{|g|+b+|h|} f \smile ((h \circ_{j-f} g) \circ_{k-f} b), & f \leq j \leq k-g \leq |h|+|f|, \\ \Gamma'_{i,i+|f|,k} &= (-1)^{|b|+|g|} (h \circ_{i-1} (f \smile g)) \circ_k b, & 1 \leq i \leq k-|f|-g \leq |h|, \\ \Gamma''_{ij,j+|g|} &= (-1)^{|b|} (h \circ_{i-1} f) \circ_{j-1} (g \smile b), & 1 \leq i \leq j-f \leq |h|, \\ \Gamma'''_{ij,|h|+f+g} &= (-1)^b ((h \circ_{i-1} f) \circ_{j-1} g) \smile b, & 1 \leq i \leq j-f \leq |h|, \end{aligned}$$

*by definition for the six edge values*

$$\begin{aligned} \Gamma_{0j,|h|+|f|+g} &\quad (f \leq j \leq |h|+|f|), & \Gamma'_{1fk} &\quad (f+g \leq k \leq |h|+|f|+|g|), \\ \Gamma'_{i,i+|f|,|h|+|f|+g} &\quad (1 \leq i \leq |h|), & \Gamma''_{1j,j+|g|} &\quad (f+1 \leq j \leq |h|+|f|), \\ \Gamma''_{i,i+f,i+f+|g|} &\quad (2 \leq i \leq |h|-1), & \Gamma'''_{i,|h|+f,h+|f|+|g|} &\quad (1 \leq i \leq |h|). \end{aligned}$$

*Proof.* See Appendix C.  $\square$

**Proof of the Main Theorem (Gerstenhaber's method).** First note that

$$\begin{aligned} \{h, f, \delta g, b\} &= \sum_{i=0}^{|h|-2} \sum_{j=i+f}^{|h|+|f|-1} \sum_{k=j+\delta g}^{|h|+|f|+|\delta g|} ((h \circ_i f) \circ_j \delta g) \circ_k b \\ &= \sum_{i=0}^{|h|-2} \sum_{j=i+f}^{|h|+|f|-1} \sum_{k=j+g+1}^{|h|+|f|+|g|+1} ((h \circ_i f) \circ_j \delta g) \circ_k b \\ &= \sum_{i=0}^{|h|-2} \sum_{j=i+f}^{|h|+|f|-1} \sum_{k=j+g}^{|h|+|f|+|g|} ((h \circ_i f) \circ_j \delta g) \circ_{k+1} b \\ &= \sum_{(i,j,k) \in T} ((h \circ_i f) \circ_j \delta g) \circ_{k+1} b \end{aligned}$$

and

$$\begin{aligned}
\{h, \delta f, g, b\} &= \sum_{i=0}^{|h|-2} \sum_{j=i+\delta f}^{|h|+|\delta f|-1} \sum_{k=j+g}^{|h|+|\delta f|+|g|} ((h \circ_i \delta f) \circ_j g) \circ_k b \\
&= \sum_{i=0}^{|h|-2} \sum_{j=i+f+1}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|+1} ((h \circ_i \delta f) \circ_j g) \circ_k b \\
&= \sum_{i=0}^{|h|-2} \sum_{j=i+f}^{|h|+|f|-1} \sum_{k=j+g}^{|h|+|f|+|g|} ((h \circ_i \delta f) \circ_{j+1} g) \circ_{k+1} b \\
&= \sum_{(i,j,k) \in T} ((h \circ_i \delta f) \circ_{j+1} g) \circ_{k+1} b.
\end{aligned}$$

By using Lemma 6.5 we have

$$\begin{aligned}
&\delta\{h, f, g, b\} - \{h, f, g, \delta b\} - (-1)^{|b|}\{h, f, \delta g, b\} - (-1)^{|b|+|g|}\{h, \delta f, g, b\} \\
&= \sum_{(i,j,k) \in T} (\Gamma_{i+1,j+1,k+1} + \Gamma'_{i+1,j+1,k+1} + \Gamma''_{i+1,j+1,k+1} + \Gamma'''_{i+1,j+1,k+1}) \\
&= \sum_{(i,j,k) \in T'} (\Gamma_{ijk} + \Gamma'_{ijk} + \Gamma''_{ijk} + \Gamma'''_{ijk}).
\end{aligned}$$

Now use Lemma 6.7 to see that

$$\begin{aligned}
(-1)^{|f|+|g|+|b|}\{\delta h, f, g, b\} &= (-1)^{|f|+|g|+|b|} \sum_{i=0}^{|\delta h|-2} \sum_{j=i+f}^{|\delta h|+|f|-1} \sum_{k=j+g}^{|\delta h|+|f|+|g|} ((\delta h \circ_i f) \circ_j g) \circ_k b \\
&= \sum_{i=0}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} (\Gamma_{ijk} + \Gamma'_{i+1,j,k} + \Gamma''_{i+1,j+1,k} + \Gamma'''_{i+1,j+1,k+1}) \\
&= \sum_{i=0}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma_{ijk} + \sum_{i=1}^{|h|} \sum_{j=i+|f|}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma'_{ijk} \\
&\quad + \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+|g|}^{|h|+|f|+|g|} \Gamma''_{ijk} + \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma'''_{ijk} \\
&= \sum_{i=1}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma_{ijk} + \sum_{j=f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma_{0jk} \\
&\quad + \sum_{i=1}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma'_{ijk} + \sum_{i=1}^{|h|} \sum_{k=i+|f|+g}^{|h|+|f|+|g|} \Gamma'_{i,i+|f|,k} \\
&\quad + \sum_{i=1}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma''_{ijk} + \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+|f|} \Gamma''_{ij,j+|g|}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i=1}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma'''_{ijk} + \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+f} \Gamma'''_{ij,h+f+|g|} \\
& = \sum_{(i,j,k) \in T'} (\Gamma_{ijk} + \Gamma'_{ijk} + \Gamma''_{ijk} + \Gamma'''_{ijk}) \\
& \quad \boxed{
\begin{aligned}
& + \sum_{j=f}^{|h|+|f|} \sum_{k=j+g}^{|h|+|f|+|g|} \Gamma_{0jk} + \sum_{i=1}^{|h|} \sum_{k=i+|f|+g}^{|h|+|f|+|g|} \Gamma'_{i,i+|f|,k} \\
& + \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+f} \Gamma''_{ij,j+|g|} + \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+f} \Gamma''_{ij,h+f+|g|}
\end{aligned}
}
\end{aligned}$$

One can see that the resulting  $\boxed{\text{boxed formula}}$  is a sum over the *boundary*  $\partial T'_{env}$  of the truncated envelope  $T'_{env}$  of  $T'$ . By cancelling the sums  $\sum_{T'}$  and using the Boundary Lemma, we finally obtain

$$\begin{aligned}
\text{dev}_{\{.,.,.\}} \delta(h \otimes f \otimes g \otimes b) & = (-1)^{|g|+|b|+|h|f} \sum_{j=f}^{|f|+|h|} \sum_{k=j+g}^{|h|+|f|+|g|} f \smile ((h \circ_{j-f} g) \circ_{k-f} b) \\
& - (-1)^{|g|+|b|} \sum_{i=1}^{|h|} \sum_{k=i+|f|+g}^{|h|+|f|+|g|} (h \circ_{i-1} (f \smile g)) \circ_k b \\
& - (-1)^{|b|} \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+f} (h \circ_{i-1} f) \circ_{j-1} (g \smile b) \\
& - (-1)^{|b|} \sum_{i=1}^{|h|} \sum_{j=i+f}^{|h|+f} ((h \circ_{i-1} f) \circ_{j-1} g) \smile b \\
& = (-1)^{|b|} \left[ (-1)^{|h|f+|g|} \sum_{j=0}^{|h|-1} \sum_{k=j+g}^{|h|+|g|} f \smile ((h \circ_j g) \circ_k b) \right. \\
& \quad - (-1)^{|g|} \sum_{i=0}^{|h|-1} \sum_{j=i+f+g}^{|h|+|f|+|g|} (h \circ_i (f \smile g)) \circ_j b \\
& \quad \left. - \sum_{i=0}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} (h \circ_i f) \circ_j (g \smile b) + \sum_{i=0}^{|h|-1} \sum_{j=i+f}^{|h|+|f|} ((h \circ_i f) \circ_j g) \smile b \right] \\
& = (-1)^{|b|} \left[ (-1)^{|h|f+|g|} f \smile \{h, g, b\} - (-1)^{|g|} \{h, f \smile g, b\} - \{h, f, g \smile b\} \right. \\
& \quad \left. + \{h, f, g\} \smile b \right],
\end{aligned}$$

which is the required formula.  $\square$

## 7. APPENDIX A

**Proof of Lemma 6.5.** First note that

$$\begin{aligned} -\delta_\mu(((h \circ_i f) \circ_j g) \circ_k b) &= (-1)^{|h|+|f|+|g|+|b|} \mathbf{I} \smile (((h \circ_i f) \circ_j g) \circ_k b) \\ &\quad + \sum_{s=0}^{|h|+|f|+|g|+|b|} (((h \circ_i f) \circ_j g) \circ_k b) \circ_s \mu + (((h \circ_i f) \circ_j g) \circ_k b) \smile \mathbf{I}. \end{aligned}$$

By using the composition relations

$$(((h \circ_i f) \circ_j g) \circ_k b) \circ_s \mu = \begin{cases} (-1)^{|b|} (((h \circ_i f) \circ_j g) \circ_s \mu) \circ_{k+1} b, & \text{if } 0 \leq s \leq k-1 \\ ((h \circ_i f) \circ_j g) \circ_k (b \circ_{s-k} \mu), & \text{if } k \leq s \leq k+|b| \\ (-1)^{|b|} (((h \circ_i f) \circ_j g) \circ_{s-|b|} \mu) \circ_k b, & \text{if } k+b \leq s \end{cases}$$

cut the above sum  $\sum_{s=0}^{|h|+|f|+|g|+|b|}$  in three pieces,

$$\begin{aligned} -\delta(((h \circ_i f) \circ_j g) \circ_k b) &= (-1)^{|h|+|f|+|g|+|b|} \mathbf{I} \smile (((h \circ_i f) \circ_j g) \circ_k b) \\ &\quad + (-1)^{|b|} \sum_{s=0}^{k-1} (((h \circ_i f) \circ_j g) \circ_s \mu) \circ_{k+1} b + \sum_{s=k}^{k+|b|} ((h \circ_i f) \circ_j g) \circ_k (b \circ_{s-k} \mu) \\ &\quad + (-1)^{|b|} \sum_{s=k+b}^{|h|+|f|+|g|+|b|} (((h \circ_i f) \circ_j g) \circ_{s-|b|} \mu) \circ_k b + (((h \circ_i f) \circ_j g) \circ_k b) \smile \mathbf{I}. \end{aligned}$$

Next, cut the first sum  $\sum_{s=0}^{k-1}$  in three pieces by using the composition relations

$$((h \circ_i f) \circ_j g) \circ_s \mu = \begin{cases} (-1)^{|g|} ((h \circ_i f) \circ_s \mu) \circ_{j+1} g, & \text{if } 0 \leq s \leq j-1 \\ (h \circ_i f) \circ_j (g \circ_{s-j} \mu), & \text{if } j \leq s \leq j+|g| \\ (-1)^{|g|} ((h \circ_i f) \circ_{s-|g|} \mu) \circ_j g, & \text{if } j+g \leq s \leq k-1, \end{cases}$$

and obtain

$$\begin{aligned} -\delta(((h \circ_i f) \circ_j g) \circ_k b) &= (-1)^{|h|+|f|+|g|+|b|} \mathbf{I} \smile (((h \circ_i f) \circ_j g) \circ_k b) \\ &\quad + (-1)^{|b|+|g|} \sum_{s=0}^{j-1} (((h \circ_i f) \circ_s \mu) \circ_{j+1} g) \circ_{k+1} b \\ &\quad + (-1)^{|b|} \sum_{s=j}^{j+|g|} ((h \circ_i f) \circ_j (g \circ_{s-j} \mu)) \circ_{k+1} b \\ &\quad + (-1)^{|b|+|g|} \sum_{s=j+g}^{k-1} (((h \circ_i f) \circ_{s-|g|} \mu) \circ_j g) \circ_{k+1} b \\ &\quad + \sum_{s=k}^{k+|b|} ((h \circ_i f) \circ_j g) \circ_k (b \circ_{s-k} \mu) \\ &\quad + (-1)^{|b|} \sum_{s=k+b}^{|h|+|f|+|g|+|b|} (((h \circ_i f) \circ_j g) \circ_{s-|b|} \mu) \circ_k b \\ &\quad + (((h \circ_i f) \circ_j g) \circ_k b) \smile \mathbf{I}. \end{aligned}$$

Next, cut the first sum  $\sum_{s=0}^{j-1}$  in three pieces by using the composition relations

$$(h \circ_i f) \circ_s \mu = \begin{cases} (-1)^{|f|} (h \circ_s \mu) \circ_{i+1} f, & \text{if } 0 \leq s \leq i-1 \\ h \circ_i (f \circ_{s-i} \mu), & \text{if } i \leq s \leq i+|f| \\ (-1)^{|f|} (h \circ_{s-|f|} \mu) \circ_i f, & \text{if } i+f \leq s \leq j-1, \end{cases}$$

and desuspend summation ranges,

$$\sum_{s=n}^{n+|x|} x \circ_{s-n} \mu = \sum_{s=0}^{|x|} x \circ_s \mu = x \bullet \mu = -(-1)^{|x|} \mathbf{I} \smile x - \delta_\mu x - x \smile \mathbf{I}$$

for the pairs  $(n, x) = (k, b)$ ,  $(j, g)$  and  $(i, f)$ ,

$$\begin{aligned} \sum_{s=i+f}^{j-1} h \circ_{s-|f|} \mu &= \sum_{s=i+1}^{j-f} h \circ_s \mu, \\ \sum_{s=j+g}^{k-1} (h \circ_i f) \circ_{s-|g|} \mu &= (-1)^{|f|} \sum_{s=j+g}^{k-1} (h \circ_{s-|g|-|f|} \mu) \circ_i f = (-1)^{|f|} \sum_{s=j-|f|+1}^{k-|f|-g} (h \circ_s \mu) \circ_i f, \\ \sum_{s=k+b}^{|h|+|f|+|g|+|b|} ((h \circ_i f) \circ_j g) \circ_{s-|b|} \mu &= (-1)^{|g|} \sum_{s=k+b}^{|h|+|f|+|g|+|b|} ((h \circ_i f) \circ_{s-|b|-|g|} \mu) \circ_j g \\ &= (-1)^{|f|+|g|} \sum_{s=k+b}^{|h|+|f|+|g|+|b|} ((h \circ_{s-|b|-|g|-|f|} \mu) \circ_i f) \circ_j g \\ &= (-1)^{|f|+|g|} \sum_{s=k-|f|-|g|+1}^{|h|} ((h \circ_s \mu) \circ_i f) \circ_j g \end{aligned}$$

to obtain the required formula.  $\square$

## 8. APPENDIX B

**8.1. Proof of Lemma 6.7.** First note that

$$\begin{aligned} \Gamma_{ijk} + \Gamma'_{i+1,j,k} + \Gamma''_{i+1,j+1,k} + \Gamma'''_{i+1,j+1,k+1} \\ = -(-1)^{|h|+|f|+|g|+|b|} \mathbf{I} \smile (((h \circ_{i-1} f) \circ_{j-1} g) \circ_{k-1} b) \\ - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\ + (-1)^{|f|+|g|+|b|} ((h \circ_{i-1} (\mathbf{I} \smile f)) \circ_j g) \circ_k b \\ + (-1)^{|g|+|b|} ((h \circ_i (f \smile \mathbf{I})) \circ_j g) \circ_k b \\ - (-1)^{|f|+|g|+|b|} \sum_{s=i+1}^{j-f-1} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\ + (-1)^{|g|+|b|} ((h \circ_i f) \circ_{j-1} (\mathbf{I} \smile g)) \circ_k b \end{aligned}$$

$$\begin{aligned}
& + (-1)^{|b|}((h \circ_i f) \circ_j (g \smile I)) \circ_k b \\
& - (-1)^{|f|+|g|+|b|} \sum_{s=j-|f|+1}^{k-f-g} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\
& + (-1)^{|b|}((h \circ_i f) \circ_j g) \circ_{k-1} (I \smile b) \\
& + ((h \circ_i f) \circ_j g) \circ_k (b \smile I) \\
& - (-1)^{|f|+|g|+|b|} \sum_{s=k-|f|-|g|+1}^{|h|} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\
& - (((h \circ_i f) \circ_j g) \circ_k b) \smile I.
\end{aligned}$$

We must compare it term by term with

$$\begin{aligned}
-((\delta h \circ_i f) \circ_j g) \circ_k b &= \left( \left( (-1)^{|h|} I \smile h + \sum_{s=0}^{|h|} h \circ_s \mu + h \smile I \right) \circ_i f \right) \circ_j g \circ_k b \\
&= (-1)^{|h|}((I \smile h) \circ_i f) \circ_j g \circ_k b + \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\
&\quad + (((h \circ_{i-1} \mu) \circ_i f) \circ_j g) \circ_k b + (((h \circ_i \mu) \circ_i f) \circ_j g) \circ_k b \\
&\quad + \sum_{s=i+1}^{j-f-1} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b + (((h \circ_{j-f} \mu) \circ_i f) \circ_j g) \circ_k b \\
&\quad + (((h \circ_{j-|f|} \mu) \circ_i f) \circ_j g) \circ_k g + \sum_{s=j-|f|+1}^{k-f-g} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b \\
&\quad + (((h \circ_{k-|f|-g} \mu) \circ_i f) \circ_j g) \circ_k b + (((h \circ_{k-|f|-|g|} \mu) \circ_i f) \circ_j g) \circ_k b \\
&\quad + \sum_{s=k-|f|-|g|+1}^{|h|} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b + (((h \smile I) \circ_i f) \circ_j g) \circ_k b.
\end{aligned}$$

Now, recall the sign  $(-1)^{|f|+|g|+|b|}$  and use composition relations to note the **ground identities**

$$\begin{aligned}
((I \smile h) \circ_i f) \circ_j g \circ_k b &= (I \smile (h \circ_{i-1} f)) \circ_j g \circ_k b \\
&= (I \smile ((h \circ_{i-1} f) \circ_{j-1} g)) \circ_k b \\
&= I \smile (((h \circ_{i-1} f) \circ_{j-1} g) \circ_{k-1} b), \\
(h \circ_{i-1} \mu) \circ_i f &= h \circ_{i-1} (\mu \circ_1 f) = -h \circ_{i-1} (I \smile f), \\
(h \circ_i \mu) \circ_i f &= h \circ_i (\mu \circ_0 f) = (-1)^f h \circ_i (f \smile I), \\
((h \circ_{j-f} \mu) \circ_i f) \circ_j g &= (-1)^{|f|}((h \circ_i f) \circ_{j-1} \mu) \circ_j g \\
&= (-1)^{|f|}(h \circ_i f) \circ_{j-1} (\mu \circ_1 g) \\
&= (-1)^f (h \circ_i f) \circ_{j-1} (I \smile g), \\
((h \circ_{j-|f|} \mu) \circ_i f) \circ_j g &= (-1)^{|f|}((h \circ_i f) \circ_j \mu) \circ_j g
\end{aligned}$$

$$\begin{aligned}
&= (-1)^{|f|} (h \circ_i f) \circ_j (\mu \circ_0 g) \\
&= (-1)^{|f|+g} (h \circ_i f) \circ_j (g \smile \mathbf{I}), \\
((h \circ_{k-|f|-g} \mu) \circ_i f) \circ_j g) \circ_k b &= (-1)^{|f|} (((h \circ_i f) \circ_{k-g} \mu) \circ_j g) \circ_k b \\
&= (-1)^{|f|+|g|} (((h \circ_i f) \circ_j g) \circ_{k-1} \mu) \circ_k b \\
&= (-1)^{|f|+|g|} ((h \circ_i f) \circ_j g) \circ_{k-1} (\mu \circ_1 b) \\
&= (-1)^{|f|+g} ((h \circ_i f) \circ_j g) \circ_{k-1} (\mathbf{I} \smile b), \\
((h \circ_{k-|f|-|g|} \mu) \circ_i f) \circ_j g) \circ_k b &= (-1)^{|f|} (((h \circ_i f) \circ_{k-|g|} \mu) \circ_j g) \circ_k b \\
&= (-1)^{|f|+|g|} ((h \circ_i f) \circ_j g) \circ_k \mu \circ_k b \\
&= (-1)^{|f|+|g|} ((h \circ_i f) \circ_j g) \circ_k (\mu \circ_0 b) \\
&= (-1)^{|f|+|g|+b} ((h \circ_i f) \circ_j g) \circ_k (b \smile \mathbf{I}), \\
(((h \smile \mathbf{I}) \circ_i f) \circ_j g) \circ_k b &= (-1)^{|f|} (((h \circ_i f) \smile \mathbf{I}) \circ_j g) \circ_k b \\
&= (-1)^{|f|+|g|} (((h \circ_i f) \circ_j g) \smile \mathbf{I}) \circ_k b \\
&= (-1)^{|f|+|g|+|b|} (((h \circ_i f) \circ_j g) \circ_k b) \smile \mathbf{I},
\end{aligned}$$

which prove the required formula.  $\square$

**8.2. Proposition/recapitulation.** In a pre-operad  $C$ , for  $(i, j, k) \in T'$  the auxiliary variables read

$$\begin{aligned}
\Gamma_{ijk} &:= -(-1)^{|h|+|f|+|g|+|b|} ((\mathbf{I} \smile h) \circ_i f) \circ_j g) \circ_k b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{i-1} (((h \circ_s \mu) \circ_i f) \circ_j g) \circ_k b, \\
\Gamma'_{ijk} &:= -(-1)^{|f|+|g|+|b|} \sum_{s=i-1}^{j-f} (((h \circ_s \mu) \circ_{i-1} f) \circ_j g) \circ_k b, \\
\Gamma''_{ijk} &:= -(-1)^{|f|+|g|+|b|} \sum_{s=j-f}^{k-f-|g|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_k b, \\
\Gamma'''_{ijk} &:= -(-1)^{|f|+|g|+|b|} \sum_{s=k-f-|g|}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{k-1} b \\
&\quad - (-1)^{|f|+|g|+|b|} (((h \smile \mathbf{I}) \circ_{i-1} f) \circ_{j-1} g) \circ_{k-1} b.
\end{aligned}$$

*Proof.* Use the **ground identities** from the previous section 8.1.  $\square$

## 9. APPENDIX C

**Proof of the Boundary Lemma.** We calculate the boundary values of the auxiliary variables in a standard way, by using Lemma 6.7 and recapitulation formulae 8.2. First prove that

$$\Gamma''_{ij,j+|g|} = (-1)^{|b|} (h \circ_{i-1} f) \circ_{j-1} (g \smile b)$$

for  $2 \leq i \leq |h| - 2$ ,  $i + f + 1 \leq j \leq |h| + |f|$ . Use Lemma 6.7 and recapitulation formulae 8.2 to note that

$$\begin{aligned}
& \Gamma_{i-1,j-1,j+|g|} + \Gamma'_{i,j-1,j+|g|} + \Gamma''_{ij,j+|g|} + \Gamma'''_{ij,j+g} \\
&= -(-1)^{|h|+|f|+|g|+|b|} (((I \smile h) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=i-1}^{j-f-1} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b + \Gamma''_{ij,j+|g|} \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=j-|f|}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
&\quad - (-1)^{|f|+|g|+|b|} (((h \smile I) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b.
\end{aligned}$$

We must compare it term by term with

$$\begin{aligned}
& -((\delta h \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b = (-1)^{|h|} (((I \smile h) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
& \quad + \sum_{s=0}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b + (((h \smile I) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
&= (-1)^{|h|} (((I \smile h) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b + \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
& \quad + \sum_{s=i-1}^{j-f-1} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b + (((h \circ_{j-f} \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
& \quad + \sum_{s=j-|f|}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b + (((h \smile I) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b.
\end{aligned}$$

Now recall the sign  $(-1)^{|f|+|g|+|b|}$  and use composition relations to note that

$$\begin{aligned}
& (((h \circ_{j-f} \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b = (-1)^{|f|} (((h \circ_{i-1} f) \circ_{j-1} \mu) \circ_{j-1} g) \circ_{j+|g|} b \\
&= (-1)^{|f|} ((h \circ_{i-1} f) \circ_{j-1} (\mu \circ_0 g)) \circ_{j+|g|} b \\
&= (-1)^{|f|} (h \circ_{i-1} f) \circ_{j-1} ((\mu \circ_0 g) \circ_g b) \\
&= (-1)^{|f|+g} (h \circ_{i-1} f) \circ_{j-1} (g \smile b),
\end{aligned}$$

which lead one to the required formula for  $\Gamma''_{ij,j+|g|}$ . Also note that the last identities hold if  $1 \leq i \leq |h|$ ,  $i + f \leq j \leq |h| + f$ . The latter inequalities represent a *projection* on the  $ij$ -coordinate plane of the following face of  $T'_{env}$ :

$$1 \leq i \leq |h|, \quad i + f \leq j \leq |h| + f, \quad k = j + |g|.$$

Therefore, in this case (projection) we have

$$\Gamma''_{ij,j+|g|} = (-1)^{|b|} (h \circ_{i-1} f) \circ_{j-1} (g \smile b)$$

$$\begin{aligned}
&= -(-1)^{|f|+|g|+|b|}(((h \circ_{j-f} \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b \\
&= -(-1)^{|f|+|g|+|b|} \sum_{s=j-f}^{j-f} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{j+|g|} b
\end{aligned}$$

and the recapitulation formula

$$\Gamma''_{ijk} = -(-1)^{|f|+|g|+|b|} \sum_{s=j-f}^{k-f-|g|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_k b$$

holds in the tetrahedron

$$1 \leq i \leq |h|, \quad i + f \leq j \leq |h| + f, \quad j + |g| \leq k \leq |h| + f + |g|.$$

Next prove that

$$\Gamma'_{i,i+|f|,k} = (-1)^{|b|+|g|}(h \circ_{i-1} (f \smile g)) \circ_k b$$

for  $2 \leq i \leq |h| - 1$ ,  $i + |f| + g \leq k \leq |h| + |f| + |g|$ . Use Lemma 6.7 and recapitulation formulae 8.2 to note that

$$\begin{aligned}
&\Gamma_{i-1,i+|f|,k} + \Gamma'_{i,i+|f|,k} + \Gamma''_{i,i+f,k} + \Gamma'''_{i,i+f,k+1} \\
&= -(-1)^{|h|+|f|+|g|+|b|} (((I \smile h) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b + \Gamma'_{i,i+|f|,k} \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=i}^{k-f-|g|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&\quad - (-1)^{|f|+|g|+|b|} \sum_{s=k-|f|-|g|}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&\quad - (-1)^{|f|+|g|+|b|} (((h \smile I) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b.
\end{aligned}$$

We have to compare it term by term with

$$\begin{aligned}
&-((\delta h \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b = (-1)^{|h|} (((I \smile h) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&\quad + \sum_{s=0}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b + (((h \smile I) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&= (-1)^{|h|} (((I \smile h) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b + \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&\quad + (((h \circ_{i-1} \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b + \sum_{s=i}^{k-f-|g|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\
&\quad + \sum_{s=k-|f|-|g|}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b + (((h \smile I) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b.
\end{aligned}$$

Now recall the sign  $(-1)^{|f|+|g|+|b|}$  and use composition relations to note that

$$\begin{aligned} ((h \circ_{i-1} \mu) \circ_{i-1} f) \circ_{i+|f|} g &= (h \circ_{i-1} (\mu \circ_0 f)) \circ_{i-1+|f|} g \\ &= h \circ_{i-1} ((\mu \circ_0 f) \circ_f g) \\ &= (-1)^f h \circ_{i-1} (f \smile g), \end{aligned}$$

which lead one to the required formula for  $\Gamma'_{i,i+|f|,k}$ . Also note that the last identities hold if  $1 \leq i \leq |h|$ ,  $i + |f| + g \leq k \leq |h| + |f| + g$ . The latter inequalities represent a *projection* on the  $ik$ -coordinate plane of the following face of  $T'_{env}$ :

$$1 \leq i \leq |h|, \quad j = i + |f|, \quad i + |f| + g \leq k \leq |h| + |f| + g.$$

Therefore, in this case (projection) we have

$$\begin{aligned} \Gamma'_{i,i+|f|,k} &= (-1)^{|b|+|g|} (h \circ_{i-1} (f \smile g)) \circ_k b \\ &= -(-1)^{|f|+|g|+|b|} (((h \circ_{i-1} \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \\ &= -(-1)^{|f|+|g|+|b|} \sum_{s=i-1}^{i-1} (((h \circ_s \mu) \circ_{i-1} f) \circ_{i+|f|} g) \circ_k b \end{aligned}$$

and the recapitulation formula

$$\Gamma'_{ijk} = -(-1)^{|f|+|g|+|b|} \sum_{s=i-1}^{j-f} (((h \circ_s \mu) \circ_{i-1} f) \circ_j g) \circ_k b$$

holds in the tetrahedron

$$1 \leq i \leq |h|, \quad i + |f| \leq j \leq |h| + |f|, \quad j + g \leq k \leq |h| + |f| + g.$$

Next prove that

$$\Gamma_{0jk} = (-1)^{|g|+|b|+|h|f} f \smile ((h \circ_{j-f} g) \circ_{k-f} b)$$

for  $f \leq j \leq |h| + |f| - 1$ ,  $j + g \leq k \leq |h| + |f| + |g|$ . Use Lemma 6.7 and recapitulation formulae 8.2 to note that

$$\begin{aligned} \Gamma_{0jk} + \Gamma'_{1jk} + \Gamma''_{1,j+1,k} + \Gamma'''_{1,j+1,k+1} &= \Gamma_{0jk} \\ &- (-1)^{|f|+|g|+|b|} \sum_{s=0}^{j-f} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b \\ &- (-1)^{|f|+|g|+|b|} \sum_{s=j-|f|}^{k-f-|g|} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b \\ &- (-1)^{|f|+|g|+|b|} \sum_{s=k-|f|-|g|}^{|h|} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b \\ &- (-1)^{|f|+|g|+|b|} (((h \smile \mathbf{I}) \circ_0 f) \circ_j g) \circ_k b. \end{aligned}$$

We must compare it term by term with

$$-((\delta h \circ_0 f) \circ_j g) \circ_k b = (-1)^{|h|} (((\mathbf{I} \smile h) \circ_0 f) \circ_j g) \circ_k b$$

$$\begin{aligned}
& + \sum_{s=0}^{|h|} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b + (((h \smile \mathbf{I}) \circ_0 f) \circ_j g) \circ_k b \\
& = (-1)^{|h|} (((\mathbf{I} \smile h) \circ_0 f) \circ_j g) \circ_k b + \sum_{s=0}^{j-f} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b \\
& + \sum_{s=j-|f|}^{k-f-|g|} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b \\
& + \sum_{s=k-|f|-|g|}^{|h|} (((h \circ_s \mu) \circ_0 f) \circ_j g) \circ_k b + (((h \smile \mathbf{I}) \circ_0 f) \circ_j g) \circ_k b.
\end{aligned}$$

Now recall the sign  $(-1)^{|f|+|g|+|b|}$  and use composition relations to note that

$$\begin{aligned}
(-1)^{|h|} (((\mathbf{I} \smile h) \circ_0 f) \circ_j g) \circ_k b & = -(-1)^{|h|} (((\mu \circ_1 h) \circ_0 f) \circ_j g) \circ_k b \\
& = -(-1)^{|h|+|h||f|} (((\mu \circ_0 f) \circ_f h) \circ_j g) \circ_k b \\
& = -(-1)^{f|h|} ((\mu \circ_0 f) \circ_f (h \circ_{j-f} g)) \circ_k b \\
& = -(-1)^{f|h|} (\mu \circ_0 f) \circ_f ((h \circ_{j-f} g) \circ_{k-f} b) \\
& = (-1)^{f|h|+|f|} f \smile ((h \circ_{j-f} g) \circ_{k-f} b),
\end{aligned}$$

which leads one to the required formula for  $\Gamma_{0jk}$ . Also note that the last identities hold if  $f \leq j \leq |h| + |f|$ ,  $j + g \leq k \leq |h| + |f| + g$ . The latter inequalities represent a *projection* on the  $jk$ -coordinate plane of the following face of  $T'_{env}$ :

$$i = 0, \quad f \leq j \leq |h| + |f|, \quad j + g \leq k \leq |h| + |f| + g.$$

Therefore, in this case (projection) we have

$$\begin{aligned}
\Gamma_{0jk} & = (-1)^{|g|+|b|+|h|f} f \smile ((h \circ_{j-f} g) \circ_{k-f} b) \\
& = -(-1)^{|h|+|f|+|g|+|b|} (((\mathbf{I} \smile h) \circ_0 f) \circ_j g) \circ_k b.
\end{aligned}$$

At last prove that

$$\Gamma'''_{ij,|h|+f+g} = (-1)^b ((h \circ_{i-1} f) \circ_{j-1} g) \smile b$$

for  $1 \leq i \leq |h|$ ,  $i + f \leq j \leq |h| + f$ . Use Lemma 6.7 and recapitulation formulae 8.2. If  $i = 1$ , then

$$\begin{aligned}
& \Gamma_{0,j-1,|h|+|f|+g} + \Gamma'_{1,j-1,|h|+|f|+g} + \Gamma''_{1j,|h|+|f|+g} + \Gamma'''_{1j,|h|+f+g} \\
& = -(-1)^{|h|+|f|+|g|+|b|} (((\mathbf{I} \smile h) \circ_0 f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\
& - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{j-f-1} (((h \circ_s \mu) \circ_0 f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\
& - (-1)^{|f|+|g|+|b|} \sum_{s=j-f}^{|h|} (((h \circ_s \mu) \circ_0 f) \circ_{j-1} g) \circ_{|h|+|f|+g} b + \Gamma'''_{1j,|h|+f+g} \\
& = -(-1)^{|h|+|f|+|g|+|b|} (((\mathbf{I} \smile h) \circ_0 f) \circ_{j-1} g) \circ_{|h|+|f|+g} b
\end{aligned}$$

$$-(-1)^{|f|+|g|+|b|} \sum_{s=0}^{|h|} (((h \circ_s \mu) \circ_0 f) \circ_{j-1} g) \circ_{|h|+|f|+g} b + \Gamma'''_{1j,|h|+f+g}.$$

If  $2 \leq i \leq |h|$ , then

$$\begin{aligned} & \Gamma_{i-1,j-1,|h|+|f|+g} + \Gamma'_{i,j-1,|h|+|f|+g} + \Gamma''_{ij,|h|+|f|+g} + \Gamma'''_{ij,|h|+f+g} \\ &= -(-1)^{|h|+|f|+|g|+|b|} (((I \smile h) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\ & \quad - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{i-2} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\ & \quad - (-1)^{|f|+|g|+|b|} \sum_{s=i-1}^{j-f-1} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\ & \quad - (-1)^{|f|+|g|+|b|} \sum_{s=j-f}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b + \Gamma'''_{ij,|h|+f+g} \\ &= -(-1)^{|h|+|f|+|g|+|b|} (((I \smile h) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\ & \quad - (-1)^{|f|+|g|+|b|} \sum_{s=0}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b + \Gamma'''_{ij,|h|+f+g}. \end{aligned}$$

So the both cases ( $i = 1$  and  $2 \leq i \leq |h|$ ) can be obtained from unique formula for  $1 \leq i \leq |h|$ . We must compare it term by term with

$$\begin{aligned} & ((\delta h \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b = -(-1)^{|h|} (((I \smile h) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\ & \quad - \sum_{s=0}^{|h|} (((h \circ_s \mu) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b - (((h \smile I) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b. \end{aligned}$$

Now recall the sign  $(-1)^{|f|+|g|+|b|}$  and use composition relations to note that

$$\begin{aligned} & -(((h \smile I) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b = (-1)^{|h|} (((\mu \circ_0 h) \circ_{i-1} f) \circ_{j-1} g) \circ_{|h|+|f|+g} b \\ & \quad = (-1)^{|h|} (\mu \circ_0 ((h \circ_{i-1} f) \circ_{j-1} g)) \circ_{|h|+|f|+g} b \\ & \quad = -(-1)^{|f|+|g|} ((h \circ_{i-1} f) \circ_{j-1} g) \smile b, \end{aligned}$$

which lead one to the required formula for  $\Gamma'''_{ij,|h|+f+g}$ .  $\square$

## REFERENCES

- [1] V. Coll, M. Gerstenhaber, and S. D. Schack. Universal deformation formulas and breaking symmetry. J. Pure Appl. Algebra, 1993, **90**, 201-219.
- [2] M. Gerstenhaber. The cohomology structure of an associative ring. Ann. of Math. 1963, **78**, 267-288.
- [3] M. Gerstenhaber. On the deformation theory of rings and algebras: III. Ann. of Math. 1968, **88**, 1-34.
- [4] M. Gerstenhaber, A. Giaquinto, and S. D. Schack. Algebras, bialgebras, quantum groups, and algebraic deformations. In *Deformation Theory and Quantum Groups with Applications to Mathematical Physics*, Amherst, MA, 1990. M. Gerstenhaber and J. Stasheff, Eds. Contemp. Math. 1992, **134**, 51-92.

- [5] M. Gerstenhaber and A. A. Voronov Homotopy G-algebras and moduli space operad. Int. Math. Res. Notices, 1995, **3**, 141-153; hep-th/9409063.
- [6] L. Kluge, E. Paal, and J. Stasheff. Invitation to composition. Comm. Algebra, 2000, **28**, 1405-1422; math. QA/9908083.
- [7] L. Kluge and E. Paal. On derivation deviations in an abstract pre-operad. Comm. Algebra (to be published); math.QA/0105229.
- [8] J.-L. Loday, J. Stasheff, and A. A. Voronov, Eds. *Operads: Proceedings of Renaissance Conferences*. Contemp. Math. 1997, **202**.
- [9] M. Markl. Simplex, associahedron, and cyclohedron. Contemp. Math. 1999, **227**, 235-265; alg-geom/9707009.
- [10] J. P. May. *The Geometry of Iterated Loop Spaces*. Lecture Notes in Math. 1972, **271**.
- [11] S. Shnider and S. Sternberg. *Quantum Groups from Coalgebras to Drinfeld Algebras*. Int. Press Inc.: Boston, 1994.
- [12] J. D. Stasheff. Homotopy associativity of H-spaces, I, II. Trans. Amer. Math. Soc. 1963, **108**, 275-292, 293-312.
- [13] J. Stasheff. From operads to 'physically' inspired theories. Contemp. Math. 1997, **202**, 53-81.
- [14] A. A. Voronov and M. Gerstenhaber. Higher-order operations on the Hochschild complex. Funktsional. Anal. i Prilozhen. 1995, **29**, 1-6 (in Russian).

Department of Mathematics, Tallinn Technical University  
 Ehitaajate tee 5, 19086 Tallinn, Estonia  
 e-mails: liivi.kluge.ttu@mail.ee and epaal@edu.ttu.ee